

DRUGCHECK INFORMATION CHART

A Guide to Cutoff Levels, Urine Detection Times, Signs of Use and more.

Find online at www.sassexpress.com

Drug Type	Pharmaceutical Names	Street Names	Urine Detection Times	Signs of Use
Amphetamine (AMP)	Dexedrine, Benzedrine, Dextrostat, Adderall, Biphphetamine	Speed, Bennies, Dexies, Uppers, Footballs	Cutoff level - 1000ng/ml Detection Times: 2-4 days	Dilated pupils, decreased appetite, dry mouth & nose. Bad breath, frequent lip licking, talkative, nervous. Increased heart rate & blood pressure
Methamphetamine (MET)	Desoxyn, Methedrine	Ice, Crystal, Meth, Doe	Cutoff level - 1000ng/ml Detection Times: 2-4 days	Same as amphetamines. Overdoses cause elevated body temperature.
Ecstasy (MDMA)	None	XTC, E, Adam, Blue lips, Blue Kisses, Disco Biscuits, X-ing, Doctor	Cutoff level - 500ng/ml Detection Times: 1-3 days	Same as amphetamines. May grind teeth and clench jaw. Sweating and increase in body temperature.
MDA	None	Eve, Love Pill	Detection Times: 1-3 days	Same as amphetamines. Hallucinations, increase in body temperature. Tachycardia, agitation, very talkative
Cocaine (COC)	None	Coke, Crack, Snow, Flake, Rock, Nose Candy, Powder, Blow. Speedball is cocaine mixed with heroin	Cutoff level - 300ng/ml Detection Times: 2-4 days	Dilated pupils, increased heart rate & blood pressure. Anesthesia, Euphoria, sensitive to sound & touch. Less need for food, cardio toxic
Marijuana (THC)	Marinol	Weed, Pot, Grass, Reefer, Hash, Mary Jane, Cannabis, Columbian Acapulco Gold, Ganja, Bhang Wet or Lily is marijuana sprayed with PCP	Cutoff level - 50ng/ml Detection Times: 1-2 joints: 2-3 days. Oral Ingestion: 1-5 days. Moderate smoker: 5 days. Heavy smoker (daily): 10 days. Chronic smoker - 5 joints per day: 4-18 days or as high as a month	Increased appetite, altered perception, memory loss. Bloodshot eyes, giggle frequently, trouble walking. Disorientation, increased heart rate, vasodilatation. Dry mouth & throat, decreased respiratory rate. Panic attacks, paranoia
Phencyclidine (PCP)	Sernylan	Veterinary Anesthetic Angel Dust, Hog, Killer Weed, Wet or Lily is marijuana sprayed with PCP	Cutoff level - 25ng/ml Detection Times: Up to 14 days Chronic users may be up to 30 days	Dissociative anesthesia - analgesia & amnesia. Dissociated from environment but not asleep. Elevated pulse & blood pressure, coma
Ketamine	Ketalar	Veterinary Anesthetic Kat, Special K, Vitamin K, K, Horse Tranquilizer	Cutoff level - 25ng/ml Detection Times: Up to 14 days Up to 30 days for chronic users	Analgesia, amnesia, hallucinations. Foul-tasting drip from nasal passages. Users feel they are outside of their body. Vomiting, seizures, disassociation
Gamma Hydroxybutyric Acid, (GHB)	None	GHB, Cherry Meth, Easy Lay, Jib, Georgia Home Boy, Sleep 500, Soap, Vita G, Grievous Bodily Harm, Scoop, Goop, Fantasy, Everclear, Salty Water, Liquid X	Detection Times: 1 day	Users feel euphoric then sleepy. Used as a date rape drug. Slows breathing and heart rate
Opiates (OPI 300) (OPI 2000)	Hydromorphone(Dilaudid) Codeine, Tylenol with Codeine, Morphine (Avinza Kadian, MS Contin, Oramorph SR) Hydrocodone (Vicodin) - positive with 300 cutoff	(Heroin & Morphine based drugs) Smack, H, Skag, Dope, Junk Horse, Cheese is starter heroin for kids contains acetaminophen and dextromethorphan with Mex Brown Heroin	Cutoff level - 300 ng/ml or 2000 ng/ml Detection Times: 1-2 days	Pupils constricted. Pulse & blood pressure lowered. Body temp lowered, euphoria initially and then drowsy. Respiratory depression, convulsions, constipation, coma
Oxycodone (OXY)	Percodan, Percocet, Roxicodone, Tylox, OxyContin - a time released form of Oxycodone	OXY, OC, OX, Oxycotton, Hillbilly Heroin, Killer	Cutoff level - 100ng/ml Detection Times: 1-3 days Oxycontin has longer detection times	Same as Opiates
Propoxyphene (PPX)	Darvon, Darvocet	Yellow Footballs	Cutoff level - 300ng/ml Detection Times: 3-4 days	Same as Opiates
Barbiturates (BAR)	Nembutal, Luminal, Seconal, Tuinal, Fiorinal, Neodorm, Immenoctal, Stadolorm, Amytal, Alurate, Pentothal, Mebaral	Barbs, Downers, Tranqs, Blue Dolls, Bluebirds, Rainbows, Yellows, Reds, Red Devils, M&Ms	Cutoff level - 300ng/ml Detection Times: Short acting - 1 day (Seconal, Nembutal) Long acting: 2-3 weeks (Phenobarb)	Pupils normal, but reaction to light slow, Pulse, blood pressure & respiratory rate lowered. Sedation, stupor, coma
Methadone (MTD)	Dolophine, Methadone, L-Polamidon, Physeptone	Araidone, Fizzes	Cutoff level - 300ng/ml Detection Times: 2-3 days	Same as other Opiates. Similar to heroin but less intense. Sedation analgesia
Benzodiazepines (BZD)	Librian, Valium, Ativan, Halicon Vivol, Novopoxide, Remestan, Rohypnol, Xanax, Tranxene, Restoril, Versed, Clonopin, Rivotril, Dalmane, Serax	Bennies, Rophies	Cutoff level - 300ng/ml Detection Times: Normal dose - 2-3 days Chronic Use (of 1 or more yrs) 4-6 weeks	Drowsiness, dizziness, muscle relaxation. Hangover effect after large dose. Rohypnol used as date rape drug.
Tricyclic Antidepressants (TCA)	Amitriptyline (Elavil, Endep), Cyclobenzaprine (Flexeril), Clomipramine(Anafranil), Perphenazine (Triafon), Desiprarine (Norpramin), Imipramine (Tetranil), Nortriptyline (Pamelor), Protriptyline (Vivactil), Trimipramine (Surmortil), Promazine (Sparine), Nordoxepine, Doxepin (Sinequan)		Cutoff level - 1000ng/ml Detection Times: Therapeutic dose - 3 days Extended use - 4 weeks	Elevated body temperature. Elevated pulse & blood pressure. Pupils dilated, agitation, delirious, seizures, constipation and dry, flush skin

Note: times stated are estimates. The time determined is effected by various factors, including amount and frequency of usage, weight of the individual, other medications taken, and so forth.

This information is for reference only and not for medical use. If questions, consult with a lab director or physician knowledgeable in drugs of abuse and their effect.

List updated regularly. Found online at www.sassexpress.com